


Journal Pre-proof

Remodeling of the Cortical Structural Connectome in Posttraumatic Stress Disorder: Results from the ENIGMA-PGC PTSD Consortium

Delin Sun, Gopalkumar Rakesh, Emily K. Clarke-Rubright, Courtney C. Haswell, Mark Logue, Erin N. O’Leary, Andrew S. Cotton, Hong Xie, Emily L. Dennis, Neda Jahanshad, Lauren E. Salminen, Sophia I. Thomopoulos, Faisal Rashid, Christopher R.K. Ching, Saskia B.J. Koch, Jessie L. Frijling, Laura Nawijn, Mirjam van Zuiden, Xi Zhu, Benjamin Suarez-Jimenez, Anika Sierk, Henrik Walter, Antje Manthey, Jennifer S. Stevens, Negar Fani, Sanne J.H. van Rooij, Murray Stein, Jessica Bomyea, Inga Koerte, Kyle Choi, Steven J.A. van de Werff, Robert R.J. M. Vermeiren, Julia Herzog, Lauren A.M. Lebois, Justin T. Baker, Kerry J. Ressler, Elizabeth A. Olson, Thomas Straube, Mayuresh S. Korgaonkar, Elpiniki Andrew, Ye Zhu, Gen Li, Jonathan Ipser, Anna Hudson, Matthew Peverill, Kelly Sambrook, Evan Gordon, Lee Baugh, Gina Forster, Raluca Simons, Jeffrey Simons, Vincent Magnotta, Adi Maron-Katz, Stefan du Plessis, Seth Disner, Nicholas Davenport, Dan Grupe, Jack Nitschke, Terri A. deRoon-Cassini, Jacklynn Fitzgerald, John H. Krystal, Ifat Levy, Miranda Olf, Dick J. Veltman, Li Wang, Yuval Neria, Michael D. De Bellis, Tanja Jovanovic, Judith K. Daniels, Martha Shenton, Nic J.A. van de Wee, Christian Schmahl, Milissa L. Kaufman, Isabelle M. Rosso, Scott R. Sponheim, David Bernd Hofmann, Richard A. Bryant, Kelene A. Fercho, Dan J. Stein, Sven C. Mueller, Luan Phan, Katie A. McLaughlin, Richard J. Davidson, Christine Larson, Geoffrey May, Steven M. Nelson, Chadi G. Abdallah, Hassaan Gomaa, Amit Etkin, Soraya Seedat, Ilan Harpaz-Rotem, Israel Liberzon, Xin Wang, Paul M. Thompson, Rajendra A. Morey



PII: S2451-9022(22)00047-7

DOI: <https://doi.org/10.1016/j.bpsc.2022.02.008>

Reference: BPSC 917

To appear in: *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*

Received Date: 13 September 2021

Revised Date: 10 February 2022

Accepted Date: 18 February 2022

Please cite this article as: Sun D., Rakesh G., Clarke-Rubright E.K., Haswell C.C., Logue M., O’Leary E.N., Cotton A.S., Xie H., Dennis E.L., Jahanshad N., Salminen L.E., Thomopoulos S.I., Rashid F.,

Ching C.R.K., Koch S.B.J., Frijling J.L., Nawijn L., van Zuiden M., Zhu X., Suarez-Jimenez B., Sierk A., Walter H., Manthey A., Stevens J.S., Fani N., van Rooij S.J.H., Stein M., Bomyea J., Koerte I., Choi K., van de Werff S.J.A., Vermeiren R.R.J.M., Herzog J., Lebois L.A.M., Baker J.T., Ressler K.J., Olson E.A., Straube T., Korgaonkar M.S., Andrew E., Zhu Y., Li G., Ipser J., Hudson A., Peverill M., Sambrook K., Gordon E., Baugh L., Forster G., Simons R., Simons J., Magnotta V., Maron-Katz A., du Plessis S., Disner S., Davenport N., Grupe D., Nitschke J., deRoos-Cassini T.A., Fitzgerald J., Krystal J.H., Levy I., Olf M., Veltman D.J., Wang L., Neria Y., De Bellis M.D., Jovanovic T., Daniels J.K., Shenton M., van de Wee N.J.A., Schmahl C., Kaufman M.L., Rosso I.M., Sponheim S.R., Hofmann D.B., Bryant R.A., Fercho K.A., Stein D.J., Mueller S.C., Phan L., McLaughlin K.A., Davidson R.J., Larson C., May G., Nelson S.M., Abdallah C.G., Goma H., Etkin A., Seedat S., Harpaz-Rotem I., Liberzon I., Wang X., Thompson P.M. & Morey R.A., Remodeling of the Cortical Structural Connectome in Posttraumatic Stress Disorder: Results from the ENIGMA-PGC PTSD Consortium, *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* (2022), doi: <https://doi.org/10.1016/j.bpsc.2022.02.008>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Society of Biological Psychiatry. Published by Elsevier Inc. All rights reserved.

**Remodeling of the Cortical Structural Connectome in Posttraumatic Stress Disorder:
Results from the ENIGMA-PGC PTSD Consortium**

Running Title: Network-Based Cortical Changes in PTSD

Delin Sun ^{1,2}, Gopalkumar Rakesh ^{1,2}, Emily K. Clarke-Rubright ^{1,2}, Courtney C. Haswell ^{1,2},
Mark Logue ³⁻⁶, Erin N. O'Leary ⁷, Andrew S. Cotton ⁷, Hong Xie ⁷, Emily L. Dennis ⁸⁻¹¹, Neda
Jahanshad ⁹, Lauren E. Salminen ⁹, Sophia I. Thomopoulos ⁹, Faisal Rashid ⁹, Christopher R. K.
Ching ⁹, Saskia B. J. Koch ^{12,13}, Jessie L. Frijling ¹², Laura Nawijn ^{12,14}, Mirjam van Zuiden ¹², Xi
Zhu ^{15,16}, Benjamin Suarez-Jimenez ^{15,16}, Anika Sierk ¹⁷, Henrik Walter ¹⁷, Antje Manthey ¹⁷,
Jennifer S. Stevens ¹⁸, Negar Fani ¹⁸, Sanne J.H. van Rooij ¹⁸, Murray Stein ¹⁹, Jessica
Bomyea ¹⁹, Inga Koerte ^{8,20}, Kyle Choi ²¹, Steven J.A. van de Werff ^{22,23}, Robert R. J. M.
Vermeiren ²², Julia Herzog ²⁴, Lauren A.M. Lebois ^{25,26}, Justin T. Baker ²⁷, Kerry J. Ressler ^{18,25,}
²⁶, Elizabeth A. Olson ^{25,28}, Thomas Straube ²⁹, Mayuresh S. Korgaonkar ³⁰, Elpiniki Andrew ³¹,
Ye Zhu ^{32,33}, Gen Li ^{32,33}, Jonathan Ipser ³⁴, Anna Hudson ³⁵, Matthew Peverill ³⁶, Kelly
Sambrook ³⁷, Evan Gordon ³⁸⁻⁴⁰, Lee Baugh ⁴¹⁻⁴³, Gina Forster ^{41,42,44}, Raluca Simons ^{42,45},
Jeffrey Simons ^{43,45}, Vincent Magnotta ⁴⁶, Adi Maron-Katz ⁴⁷, Stefan du Plessis ⁴⁸, Seth Disner ^{49,}
⁵⁰, Nicholas Davenport ^{49,50}, Dan Grupe ⁵¹, Jack Nitschke ⁵², Terri A. deRoon-Cassini ⁵³,
Jacklynn Fitzgerald ⁵⁴, John H. Krystal ^{55,56}, Ifat Levy ^{55,56}, Miranda Olf ^{12,57}, Dick J. Veltman ⁵⁸,
Li Wang ^{32,33}, Yuval Neria ^{15,16}, Michael D. De Bellis ⁵⁹, Tanja Jovanovic ^{18,60}, Judith K. Daniels
⁶¹, Martha Shenton ^{8,62}, Nic J.A. van de Wee ^{22,23}, Christian Schmahl ²⁴, Milissa L. Kaufman ^{25,63},
Isabelle M. Rosso ^{25,28}, Scott R. Sponheim ^{49,50}, David Bernd Hofmann ²⁹, Richard A. Bryant ⁶⁴,
Kelene A. Fercho ^{41-43,65}, Dan J. Stein ³⁴, Sven C. Mueller ^{35,66}, Luan Phan ^{67,68}, Katie A.
McLaughlin ⁶⁹, Richard J. Davidson ^{51,52,70}, Christine Larson ⁷¹, Geoffrey May ^{38-40,72}, Steven M.
Nelson ^{38-40,72}, Chadi G. Abdallah ^{55,56}, Hassaan Gomaa ⁷³, Amit Etkin ^{47,74}, Soraya Seedat ⁴⁸,

Ilan Harpaz-Rotem^{55,56}, Israel Liberzon⁷⁵, Xin Wang⁷, Paul M. Thompson⁹, Rajendra A. Morey^{1,2*}.

¹ Brain Imaging and Analysis Center, Duke University, Durham, NC, USA.

² Department of Veteran Affairs (VA) Mid-Atlantic Mental Illness Research, Education and Clinical Center, Durham, NC, USA.

³ National Center for PTSD, VA Boston Healthcare System, Boston, MA, USA.

⁴ Department of Psychiatry, Boston University School of Medicine, Boston, MA, USA.

⁵ Biomedical Genetics, Boston University School of Medicine, Boston, MA, USA.

⁶ Department of Biostatistics, Boston University School of Public Health, Boston, MA, USA.

⁷ Department of Psychiatry, University of Toledo, Toledo, OH, USA.

⁸ Psychiatry Neuroimaging Laboratory, Brigham & Women's Hospital, Boston, MA, USA.

⁹ Imaging Genetics Center, Stevens Neuroimaging & Informatics Institute, Keck School of Medicine of USC, Marina del Rey, CA, USA.

¹⁰ Department of Neurology, University of Utah, Salt Lake City, UT, USA.

¹¹ Stanford Neurodevelopment, Affect, and Psychopathology Laboratory, Stanford, CA, USA.

¹² Department of Psychiatry, Amsterdam University Medical Centers, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands.

¹³ Donders Institute for Brain, Cognition and Behavior, Centre for Cognitive Neuroimaging, Radboud University Nijmegen, Nijmegen, The Netherlands.

¹⁴ Department of Psychiatry, Amsterdam University Medical Centers, VU University Medical Center, VU University, Amsterdam, The Netherlands.

¹⁵ Department of Psychiatry, Columbia University Medical Center, New York, NY, USA.

¹⁶ New York State Psychiatric Institute, New York, NY, USA.

¹⁷ University Medical Centre Charité, Berlin, Germany.

¹⁸ Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, GA, USA.

¹⁹ Department of Psychiatry, University of California San Diego, La Jolla, CA, USA.

²⁰ Department of Child and Adolescent Psychiatry, Psychosomatics, and Psychotherapy, Ludwig-Maximilians-Universität, Munich, Germany

²¹ Health Services Research Center, University of California, San Diego, La Jolla, CA, USA

²² Department of Psychiatry, Leiden University Medical Center, Leiden, The Netherlands.

²³ Leiden Institute for Brain and Cognition, Leiden, The Netherlands.

²⁴ Department of Psychosomatic Medicine and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, Heidelberg, Germany.

²⁵ Department of Psychiatry, Harvard Medical School, Boston, MA, USA.

²⁶ Division of Depression and Anxiety Disorders, McLean Hospital, Belmont, MA, USA.

²⁷ Institute for Technology in Psychiatry, McLean Hospital, Harvard University, Belmont, MA, USA.

²⁸ Center for Depression, Anxiety, and Stress Research, McLean Hospital, Belmont, MA, USA.

²⁹ Institute of Medical Psychology and Systems Neuroscience, University of Münster, Münster, Germany.

³⁰ Brain Dynamics Centre, Westmead Institute of Medical Research, University of Sydney, Westmead, NSW, Australia.

³¹ Department of Psychology, University of Sydney, Westmead, NSW, Australia.

³² Laboratory for Traumatic Stress Studies, Chinese Academy of Sciences Key Laboratory of Mental Health, Institute of Psychology, Chinese Academy of Sciences, Beijing, China.

³³ Department of Psychology, University of Chinese Academy of Sciences, Beijing, China.

³⁴ SA MRC Unit on Risk & Resilience in Mental Disorders, Department of Psychiatry and Neuroscience Institute, University of Cape Town, Cape Town, South Africa.

³⁵ Department of Experimental Clinical and Health Psychology, Ghent University, Ghent, Belgium.

³⁶ Department of Psychology, University of Washington, Seattle, WA, USA.

³⁷ Department of Radiology, University of Washington, Seattle, WA, USA.

³⁸ Veterans Integrated Service Network-17 Center of Excellence for Research on Returning War Veterans, Waco, TX, USA.

³⁹ Department of Psychology and Neuroscience, Baylor University, Waco, TX, USA.

- ⁴⁰ Center for Vital Longevity, School of Behavioral and Brain Sciences, University of Texas at Dallas, Dallas, TX, USA.
- ⁴¹ Division of Basic Biomedical Sciences, Sanford School of Medicine, University of South Dakota, Vermillion, SD, USA.
- ⁴² Center for Brain and Behavior Research, University of South Dakota, Vermillion, SD, USA.
- ⁴³ Sioux Falls VA Health Care System, Sioux Falls, SD, USA.
- ⁴⁴ Brain Health Research Centre, Department of Anatomy, University of Otago, Dunedin, New Zealand.
- ⁴⁵ Department of Psychology, University of South Dakota, Vermillion, SD, USA.
- ⁴⁶ Department of Radiology, Psychiatry, and Biomedical Engineering, University of Iowa, Iowa City, IA, USA.
- ⁴⁷ Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, CA, USA.
- ⁴⁸ Department of Psychiatry, Stellenbosch University, Cape Town, South Africa.
- ⁴⁹ Minneapolis VA Health Care System, Minneapolis, MN, USA.
- ⁵⁰ Department of Psychiatry, University of Minnesota, Minneapolis, MN, USA.
- ⁵¹ Center for Healthy Minds, University of Wisconsin-Madison, Madison, WI, USA.
- ⁵² Department of Psychiatry, University of Wisconsin-Madison, Madison, WI, USA.
- ⁵³ Department of Surgery, Division of Trauma and Acute Care Surgery, Medical College of Wisconsin, Milwaukee, WI, USA.
- ⁵⁴ Department of Psychology, Marquette University, Milwaukee, WI, USA.
- ⁵⁵ Division of Clinical Neuroscience, National Center for PTSD, West Haven, CT, USA.
- ⁵⁶ Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA.
- ⁵⁷ ARQ National Psychotrauma Centre, Diemen, The Netherlands.
- ⁵⁸ Department of Psychiatry, Amsterdam University Medical Centers, Amsterdam, The Netherlands.
- ⁵⁹ Healthy Childhood Brain Development Developmental Traumatology Research Program, Department of Psychiatry and Behavioral Sciences, Duke University, Durham, NC, USA.

- ⁶⁰ Department of Psychiatry and Behavioral Neuroscience, Wayne State University School of Medicine, Detroit, MI, USA.
- ⁶¹ Department of Clinical Psychology, University of Groningen, Groningen, The Netherlands.
- ⁶² VA Boston Healthcare System, Brockton Division, Brockton, MA, USA.
- ⁶³ Division of Women's Mental Health, McLean Hospital, Belmont, MA, USA.
- ⁶⁴ School of Psychology, University of New South Wales, Sydney, NSW, Australia.
- ⁶⁵ Civil Aerospace Medical Institute, US Federal Aviation Administration, Oklahoma City, OK, USA
- ⁶⁶ Department of Personality, Psychological Assessment and Treatment, University of Deusto, Bilbao, Spain.
- ⁶⁷ Department of Psychiatry, University of Illinois at Chicago, Chicago, IL, USA.
- ⁶⁸ Mental Health Service Line, Jesse Brown VA Chicago Health Care System, Chicago, IL, USA.
- ⁶⁹ Department of Psychology, Harvard University, Cambridge, MA, USA.
- ⁷⁰ Department of Psychology, University of Wisconsin-Madison, Madison, WI, USA.
- ⁷¹ Department of Psychology, University of Wisconsin-Milwaukee, Milwaukee, WI, USA.
- ⁷² Department of Psychiatry and Behavioral Science, Texas A&M University Health Science Center, Bryan, TX, USA.
- ⁷³ Department of Psychiatry, Pennsylvania State University, State College, PA, USA.
- ⁷⁴ VA Palo Alto Health Care System, Palo Alto, CA, USA.
- ⁷⁵ Department of Psychiatry, University of Michigan, Ann Arbor, MI, USA.

* Corresponding Author
Rajendra A. Morey, M.D.
40 Duke Medicine Circle, Room 414
Durham, NC 27710 USA
Phone: 919-286-0411 ext. 6425
Facsimile: 919-416-5912
E-mail: rajendra.morey@duke.edu

Abstract

Background: Posttraumatic stress disorder (PTSD) is accompanied by disrupted cortical neuroanatomy. We investigated alteration in covariance of structural networks associated with PTSD in regions that demonstrate the case-control differences in cortical thickness (CT) and surface area (SA).

Methods: Neuroimaging and clinical data were aggregated from 29 research sites in >1,300 PTSD cases and >2,000 trauma-exposed controls (age 6.2-85.2 years) by the ENIGMA-PGC PTSD working group. Cortical regions in the network were rank-ordered by effect size of PTSD-related cortical differences in CT and SA. The top- n ($n = 2$ to 148) regions with the largest effect size for PTSD > non-PTSD formed *hypertrophic networks*, the largest effect size for PTSD < non-PTSD formed *atrophic networks*, and the smallest effect size of between-group differences formed *stable networks*. The mean structural covariance (SC) of a given n -region network was the average of all positive pairwise correlations and was compared to the mean SC of 5,000 randomly generated n -region networks.

Results: Patients with PTSD, relative to non-PTSD controls, exhibited lower mean SC in CT-based and SA-based atrophic networks. Comorbid depression, sex and age modulated covariance differences of PTSD-related structural networks.

Conclusions: Covariance of structural networks based on CT and cortical SA are affected by PTSD and further modulated by comorbid depression, sex, and age. The structural covariance networks that are perturbed in PTSD comport with converging evidence from resting state functional connectivity networks and networks impacted by inflammatory processes, and stress hormones in PTSD.

Keywords: PTSD, Cortical Thickness, Surface Area, Structural Covariance, Brain Network, Depression.

Journal Pre-proof

Introduction

Posttraumatic stress disorder (PTSD) is a psychiatric condition that develops in vulnerable individuals after experiencing or witnessing a life-threatening event (1). PTSD-related changes in cortical thickness (CT) (2-5) and surface area (SA) (6, 7) are found in specific cortical regions. However, relatively little is known about how PTSD affects coordinated patterns of CT and SA differences among affected cortical regions. We sought to examine PTSD effects on networks made up of cortical regions that have the greatest and the least between-group differences in CT and SA. Identifying such networks may lend support for one or more etiopathologic models of PTSD.

Structural covariance (SC) refers to the phenomenon of covarying structural brain imaging measures between cortical regions and across individuals. This covariance may be instantiated as a structural covariance network (SCN). Structural covariance network measures are shown to be concordant with tract-based white matter connectivity, synchronous neuronal activity (e.g. functional connectivity) (8, 9), and spatial patterns of gene transcription, each of which lend biological support to SCNs (10). SCNs may index mutually trophic factors between regions that covary over the course of neurodevelopment (9). Differences in SC are associated with a variety of neuropsychiatric disorders including PTSD (11-13), schizophrenia, autism, obsessive compulsive disorder (14, 15), and even trauma exposure (16).

Our investigation of structural networks with significantly different covariance was motivated by two complementary models for understanding PTSD. **(I)** There is converging evidence that neurobiological mechanisms drive concerted patterns (covariance) of atrophy or hypertrophy across selected brain regions. There is generally more evidence supporting a role for CT-derived networks than SA-derived networks. Concerted processes operative in healthy

neurobiological states are perturbed by disease to effect patterns of network atrophy or hypertrophy. These neurobiological perturbations may manifest as changes in network covariance. Neurobiologically deleterious processes in PTSD may instigate atrophy in a coordinated manner across many regions to reveal *atrophic* networks. Deleterious processes in PTSD include chronic alteration of stress hormone levels such as cortisol and norepinephrine (17, 18), epigenetics mechanisms such as methylation (19, 20), inflammatory processes such as oxidative stress (21) and cytokines (22), and accelerated aging through the combined effect of these and other processes (23). **(II)** Alternatively, between-group differences in network SC may support one or the other prevailing neural systems models of PTSD. For instance, a dominant model of PTSD is that fear learning systems go awry in the aftermath of trauma. Behaviorally, slow or incomplete fear extinction and rapid fear-reinstatement contribute to symptoms of PTSD. Effective fear learning is dependent on the healthy function of underlying brain networks. Functional connectivity networks have been found to be congruent with structural covariance networks (24, 25). Thus, between-group differences in structural networks may simply reflect the between-group differences in functional networks, and these differences pervade networks (structural and functional) involved in fear learning behavior. It is also possible we might find hypertrophy across different networks that mediate compensatory responses to disrupted fear learning.

Wannan and colleagues (26) pioneered an innovative method to investigate the mean SC of networks constituted from regions selected by rank-ordering regions most affected by the illness of interest. This method considers only the most highly ranked regions in forming networks rather than all regions as in previous SCN analyses. Their findings in schizophrenia, suggest that some cortical networks *connecting* diverse regions may propagate cortical features from one region to another, leading to distributed cortical remodeling (9). Our approach, which modified their method, considered 3 classes of networks. (I) regions most affected by virtue of

lower CT in PTSD formed so-called *atrophic networks*. (II) Regions most affected by virtue of higher CT in PTSD formed so-called *hypertrophic networks*. (III) Regions least affected by PTSD formed *stable networks*. Rank-ordering of regions was based on the effect size of between group differences in CT or SA. The threshold for considering effect sizes (top-n) was initially set to the 2-most affected regions, and was repeated for networks of up to 148 regions (top-n = 2, 3, 4, . . . 148). Thus, networks ranging in size from 2 to 148 regions, in increments of 1 region, were tested. The SC of a network was calculated as the average effect size of the regions under consideration.

Importantly, even in the absence of statistically significant group-differences for individual cortical regions, significant group differences in covariance were detected in networks consisting of regions with the greatest between-group differences. We examined both CT-based and SA-based networks because CT and SA index distinct features of neuronal organization (27-29). This approach enhanced sensitivity to cortical morphometry and network covariance differences associated with PTSD, given that CT- and SA-based networks may reflect different interactions between regions or distinct aspects of the same interaction between regions (30, 31). Cortical volume was not examined as it is readily derived from mean CT and SA by simple multiplication of these two terms. However, CT and SA possess different biological, developmental, and genetic determinants as we discuss later.

We hypothesized that the mean covariance of n -region networks would be higher than the mean covariance of randomly selected n -region networks in both PTSD and trauma-exposed control groups. Confirmation of this hypothesis would tell us that networks constituted from selected (top- n) regions are more structurally interconnected than networks of the same size composed of randomly selected regions. We further hypothesized that mean SC would be modulated by PTSD diagnosis, as well as by PTSD and comorbid depression, given the two disorders are

highly comorbid (32). We predicted greater impact of PTSD on SA-based networks than on CT-based networks because SA generally drives performance more directly for a variety of cognitive and affective processes (33, 34). We also know that SA has an outsized role compared to CT in various neurobiological, neurodevelopmental, and neurogenetic processes. We predicted, because stable networks are made of regions that are least affected by PTSD, their covariance might be stronger than in non-PTSD since these networks of the least affected regions might compensate for disrupted networks composed of highly affected regions. We posited that because atrophic networks are made of regions most diminished by illness, the disease process would not necessarily affect all network regions in a systematic way, effectively lowering covariance. By contrast, we predicted that trauma-exposed non-PTSD subjects might be protected from developing symptoms because their atrophic networks maintained their healthy level of covariance. If hypertrophic networks result from higher-than-normal levels of trophic factors, whereas atrophic networks result from lower-than-normal levels of trophic factors, then we might reason that atrophic networks and hypertrophic networks would experience the same perturbations. However, given evidence that stress hormones and inflammatory processes play a role in regional atrophy but a lack of evidence for a role in regional hypertrophic, we predicted that hypertrophic networks would demonstrate different outcomes in relation to PTSD than atrophic networks. Specifically, we hypothesized that atrophic networks, unlike hypertrophic networks, would play a central role in modulating the effects of PTSD. Finally, we explored interaction effects of sex, age, and depression on PTSD.

Methods

Participants

All data, aggregated by the PGC-ENIGMA PTSD Working Group, was shared by 29 sites located in five countries ($N = 3,438$ for CT, and 3,436 for SA). Demographic and clinical information are summarized in **Table 1**. Only participants with clear information of PTSD diagnosis and sex were included in the following analyses (PTSD/Non-PTSD $N = 1,344/2,073$ for CT, and 1,348/2,066 for SA). The specific psychometric instruments and MRI acquisition parameters used at each study site are listed in **Supplementary Tables S1 and S2**, respectively. Detailed information of clinical measurements please see **Supplementary Methods**. All study sites obtained approval from local institutional review boards or ethics committees. All participants provided written informed consent.

Imaging Data Preprocessing

Details of imaging data preprocessing please see **Supplementary Methods**.

Harmonizing Data Across Sites

ComBat was utilized to harmonize CT and SA values by removing the effects of study sites while preserving inherent biological associations in the data (35). More details please see **Supplementary Methods**.

Adjusting for Confounding Factors

Age, age², sex, and mean whole-brain CT (or SA) estimates were regressed from the CT (or SA) estimates with a linear model (36). The age² term adjusted for possible nonlinear effects of age on CT (or SA). The mean whole-brain CT (or SA) estimate was included as a regressor to adjust for globally higher CT (or SA) estimates to reflect larger regional CT (or SA) estimates. More details please see **Supplementary Methods**.

Top-*n* Regions SC Analyses

The pipeline for the top-*n* regions SC analysis is shown in **Fig. 1A**. The top-*n* regions SC analysis was limited to networks consisting of the top-*n* ($n = 2$ to 148) cortical regions that were selected by rank-ordering PTSD-related changes in CT or SA by Cohen's *d* effect sizes (**Fig. 2** and **Supplementary Tables S4**). *Standardized* effect size estimates such as Cohen's *d* are independent of the units or magnitude of CT or SA values.

We examined three types of rank-ordering of regions to generate 3 network types (see **Fig. 1B**): (i) regions with higher CT in PTSD than non-PTSD were ordered from the largest positive to the largest negative effect size were used to construct *hypertrophic* networks, (ii) regions with higher CT in non-PTSD than PTSD were rank-ordered from the largest positive to the largest negative effect size were used to construct *atrophic* networks, and (iii) regions identified by comparing CT in PTSD to non-PTSD groups were rank-ordered from smallest to largest effect size were used to construct *stable* networks. The same approach used for CT was repeated for SA. An illustration depicting CT-based hypertrophic networks for top-3, top-10 and top-50 regions are shown in **Fig. 1C**.

Pearson correlation coefficients were computed across subjects per group between the CT (or SA) estimates for each of pairs of regions with the network. All correlation coefficients were *r*-to-

z transformed to improve normality and yielded a unique connectivity matrix for each participant group. The resulting matrix quantified the SC, which was interpreted for the present study as a measure of the connectivity strength between regions.

Actual Networks versus Random Networks

The mean SC (mean of all positive SC values within a network) of an actual network of the top- n regions was contrasted (i.e., mathematical subtraction) with the values of mean SC from 5,000 random networks consisting of n randomly chosen regions. This test was performed for SC measured in PTSD and non-PTSD groups, as well as between-group difference in SC. The randomly chosen regions were matched to the top- n regions for each value of n , based on the number of regions in each hemisphere and the mean Euclidean distance between all possible pairs of regions. The Euclidean distance was calculated based on the distance between the centers of cortical regions. This approach was conducted by generating 5,000 randomly chosen sets of n -regions that were matched on the number of regions per hemisphere. We then repeatedly replaced the set of n -regions with the largest or smallest mean distance by a randomly generated set of n -regions until the mean distance of the actual regions was not significantly different than the mean distance from the set of randomly chosen n -regions (one-sample t -test thresholded at 5%), or the number of searches exceeded 3,000.

We conducted replication analyses to test the reliability of our results, performed two tests of statistical significance that were complementary to each other – the global test and the individual test, and corrected for multiple comparisons using the false discovery rate (FDR) method (37). More details please see **Supplementary Methods**.

To test the hypothesis that brain hubs that are strongly connected with other areas (38), play a role in the spatial distribution of PTSD-related cortical changes, we investigated the association between the effect size of cortical changes for each region and the average of positive SC between said region and all the other cortical regions. Details please see **Supplementary Methods, Supplementary Results, Supplementary Discussion**.

PTSD X Sex Interaction

To investigate the modulation of sex on PTSD-related SCNs, we first divided PTSD and non-PTSD groups into male and female subgroups (see **Supplementary Table S5**). Two-way interactions were calculated by first contrasting PTSD (relative to its random networks) to non-PTSD (relative to its random networks) within each sex subgroup, and then calculating the difference between the two contrasts. More detailed comparisons between each pair of subgroups were conducted when there was a significant interaction effect between PTSD diagnosis and sex.

PTSD X Age Interaction

To investigate the modulation effect of depression on PTSD-related SCNs, we first divided PTSD and non-PTSD groups into eight decadal subgroups based on age: Age<10, 10≤Age<15, 15≤Age<20, 20≤Age<30, 30≤Age<40, 40≤Age<50, 50≤Age<60, Age≥60 (see **Supplementary Table S6**). Two-way interactions were calculated by first contrasting PTSD (relative to its random networks) to non-PTSD (relative to its random networks) within each age subgroup, and then calculating the difference between the two contrasts. More detailed comparisons between each pair of subgroups were conducted when there was a significant interaction effect between PTSD diagnosis and age.

PTSD X Depression Interaction

To investigate the modulation effect of depression on PTSD-related SCNs, we first divided PTSD and non-PTSD groups into subgroups based on depression diagnosis consisting of two subgroups: depressed and non-depressed (see **Supplementary Table S7**). Two-way interactions were calculated by first contrasting PTSD (relative to its random networks) to non-PTSD (relative to its random networks) within each depression subgroup, and then calculating the difference between the two contrasts. More detailed comparisons between each pair of subgroups were conducted when there was a significant interaction effect between PTSD diagnosis and depression.

Results

Effect Size of CT and SA differences

Effect sizes for between-group differences in CT and SA are shown in **Fig. 2** and reported in **Supplementary Tables S4**. Effect sizes ranged from -0.103 (atrophic) to +0.112 (hypertrophic) for CT, and from -0.110 (atrophic) to +0.083 (hypertrophic) for SA.

Top-n Regions SC Analyses

More detailed results of actual networks versus random networks in PTSD (**Fig. 3** and **Table 2**) and in non-PTSD (**Fig. 4** and **Table 2**) are listed in **Supplementary Results** for the methodologic confirmation.

PTSD versus Non-PTSD. As displayed in **Fig. 5** and **Table 2**, global tests showed that PTSD versus non-PTSD participants had lower mean SC in both CT-based ($p = 0.014$) and SA-based ($p = 0.024$) atrophic networks.

No significant differences in CT-based ($p = 0.098$) and SA-based ($p > 0.5$) hypertrophic networks, as well as CT-based ($p > 0.5$) and SA-based ($p > 0.5$) stable networks. No individual test results survived correction (p -values > 0.05).

Replication Analyses Results. As shown in **Fig. 6**, the global-tests results displayed in **Figs 3, 4, 5** and **Table 2** are reliable because the AUC of mean SC for the results based on all 29 sites were always located within the 95% confidence interval of the AUC of mean SC from 5,000

iterations leaving out 3 different sites with each iteration of the analysis across all types of networks.

Only a very small number of the individual-tests results were beyond their 95% confidence intervals. They are the CT-based stable network with top-24 regions in the non-PTSD group, the SA-based atrophic network with top-11 regions for the PTSD versus non-PTSD comparison, and the SA-based hypertrophic networks with top-32, 33, 34, or 35 regions for the PTSD versus non-PTSD comparison.

PTSD x Depression Interaction. As listed in **Fig. 7**, global tests showed a significant interaction effect in CT-based atrophic networks ($p = 0.029$; **Fig. 7A**). Further analyses showed that participants with depression alone had greater mean SC than the participants with PTSD and comorbid depression ($p < 0.001$), PTSD alone ($p < 0.001$), and healthy controls ($p < 0.001$).

There was a significant interaction effect in SA-based atrophic networks ($p = 0.001$; **Fig. 7B**). Further analyses showed that participants with PTSD alone had greater mean SC than participants with PTSD and comorbid depression ($p < 0.001$) and healthy controls ($p = 0.014$). Participants with depression alone also had greater mean SC than participants with PTSD and comorbid depression ($p < 0.001$) and healthy controls ($p < 0.001$).

There was a significant interaction effect in SA-based hypertrophic networks ($p = 0.014$; **Fig. 7D**). Further analyses showed that PTSD patients with co-morbid depression ($p = 0.029$) and healthy controls ($p < 0.001$) had greater mean SC than those with depression alone. No other global tests (p -values > 0.2) and no individual tests (p -values > 0.05) survived correction.

Effects of PTSD x Sex interaction. Global tests showed that females with PTSD ($p = 0.029$) and males without PTSD ($p = 0.014$) had greater mean SC in CT-based atrophic networks than females without PTSD. Males without PTSD had greater mean SC in CT-based stable networks than males with PTSD ($p = 0.014$) and females without PTSD ($p < 0.001$). No significant PTSD x sex interaction effect (global p -values > 0.1) was found in the other types of networks.

Effects of PTSD x Age interaction. An inverted-U relationship between decadal age and mean SC was observed in CT-based atrophic networks in both non-PTSD participants, peaking in the 3rd decade, and PTSD patients, peaking in the 2nd decade, and SA-based hypertrophic networks in PTSD patients and non-PTSD patients, both peaking in the 2nd decade. PTSD-related differences in mean SC were observed in different age groups, especially in the 1st decade, represented by lower mean SC in CT-based atrophic networks ($p < 0.001$) and SA-based hypertrophic networks ($p = 0.019$), as well as higher mean SC in CT-based hypertrophic ($p < 0.001$) and stable ($p < 0.001$) networks, in patients with PTSD compared to non-PTSD participants.

Discussion

We investigated CT-based and SA-based structural covariance networks composed of regions with the most atrophic, most hypertrophic, and most stable relationships to PTSD relative to trauma-exposed controls. Three network classes were composed of regions selected based on the effect size of PTSD-related differences in regional CT and SA. We compared the mean SC of these networks to random networks in PTSD and non-PTSD groups, respectively. We also investigated the role of PTSD diagnosis and PTSD severity on SC, and interaction effects of PTSD with age, sex and depression. We performed methodologic confirmation by demonstrating that PTSD and non-PTSD groups had higher SC in CT-based atrophic networks, SA-based atrophic networks, and SA-based hypertrophic networks than corresponding random networks (**Table 2** and **Fig. 3, 4**). Methodologic confirmation also showed the PTSD group had higher SC in CT-based hypertrophic networks and CT-based stable networks than corresponding random networks. Of particular interest and consistent with *a priori* hypotheses, we discovered that participants with PTSD had lower SC than trauma-exposed non-PTSD participants in CT-based and SA-based atrophic networks (**Table 2** and **Fig. 5**). Furthermore, depression alone had higher SC in both CT- and SA-based atrophic networks, and lower SC in SA-based hypertrophic networks compared to PTSD with comorbid depression and compared to healthy controls (**Fig. 7A, B, D**). Patients with PTSD alone showed lower SC in CT-based atrophic networks than patients with depression alone (**Fig. 7A**), and higher SC in SA-based atrophic networks compared to PTSD with comorbid depression and to healthy controls (**Fig. 7B**).

Our main finding shows that the networks composed of regions having the greatest PTSD-related atrophy, have significantly lower network covariance in the PTSD group than in the trauma-exposed control group. This finding was present for networks derived from both CT and

SA. A number of interpretations of this finding are tenable. First, we note a degree of consistency between CT-based and SA-based networks in our results concerned with PTSD diagnosis. Many cortical regions within networks that are affected by PTSD are strongly implicated (by definition) in PTSD such as insula, orbital frontal cortex, anterior cingulate, and subcallosal gyrus. However, our present study is not focused on the status of individual regions, but rather, in network perturbations associated with PTSD. Of particular note, the functional networks previously implicated in PTSD comport with the present structural network findings such as in low-level perceptual networks (39), salience network (40), default mode network (41), and central executive network (42), also referred to as the fronto-parietal network (43). Our finding of structural networks involving medial prefrontal cortex, posterior cingulate cortex (SA-based only), and angular gyrus, are canonical regions of the default mode network, which is also strongly implicated in PTSD. Our finding of structural networks involving anterior cingulate cortex, and insular cortex recapitulated salience network differences that have been reported in PTSD. However, our structural network findings did not recapitulate prior reports of central executive network involvement in PTSD, but the largest meta-analysis of network differences in PTSD did not find central executive network involvement (40), either. Unfortunately, there is a profound dearth of published findings on structural covariance network differences in PTSD for purposes of comparison. It is possible that the cortical networks or network mechanisms that propagate PTSD-related structural atrophy are dampened by the disease itself or dampened unevenly across brain topography. Alternatively, individuals with weaker connections in atrophic networks may be more vulnerable to PTSD. Unfortunately, our cross-sectional study design is unable to discern causal factors that contribute to PTSD.

In addition to functional networks, converging evidence of inflammatory processes, which contribute to PTSD, preferentially impact the same regions that constitute atrophic networks we identified. The medial prefrontal cortex, insula, and anterior cingulate are all preferentially

impacted by inflammatory processes that plague PTSD and other fear- and anxiety-based conditions (44). While the amygdala and hippocampus are also affected by inflammatory processes, we included only cortical structures, which have a uniquely measurable CT and SA. Stress hormones pose pronounced deleterious effects to the medial prefrontal cortex (45) and to the orbitofrontal cortex (46), which also featured prominently in the atrophic networks we linked to PTSD. Evidence of stress hormone effects on the brain are strongly informed by animal models. In humans, frontoparietal connectivity is disrupted after exposure to one month of intense academic stress (47). Thus, stress induced changes to medial prefrontal cortex, orbital frontal cortex, and frontoparietal regions were present in atrophic networks we linked to PTSD. Epigenetic effects on the brain have been linked to intergenerational trauma and its effects, particularly on the medial prefrontal cortex (48, 49). Epigenetic regulation of the FKBP5 gene in response to early trauma is implicated in PTSD pathogenesis (50). The methylation of FKBP5 CpG1 of intron 7 is associated with lower gray matter in bilateral orbital frontal gyrus (51). Epigenetic regulation at the stress-responsive genes that encode the pituitary adenylate cyclase-activating polypeptide (ADCYAP1) and CpG island methylation levels of its receptor ADCYAP1R1 predict PTSD symptom severity (50). Thus, inflammation, stress hormones and epigenetics, all appear to play a role in SC network difference linked to PTSD.

The present study extends several facets of earlier SC reports in PTSD. Broadly, the present study has three major methodological differences compared to published reports: **(i)** While we focused only on regions at the extremes of between-group differences in constructing networks, prior studies have considered all regions in such covariance networks, which compromises power compared to the feature reduction strategy we implemented. **(ii)** Our sample size (n=3,400) is 10-fold larger than any previous study (11). **(iii)** Two prior studies were focused on children and adolescents (n=88 and n=120) (12, 52) and a third study focused on remitted PTSD in adults (n=317) (11). Thus, the present study is uniquely situated with respect to

statistical power, a target population from a broad age-range, and illness chronicity. Our study extends the methodology developed by Wannan et al., (26) by investigating CT and SA of hypertrophic, atrophic, and stable networks separately rather than considering only the CT of atrophic networks. We show that some brain networks, independent of disease, mirror the spatial distribution of disease-related changes in cortical morphometry, thus confirming the work of Wannan et al. (26). Our results demonstrate for the first time that the SC of three different network classes are each uniquely associated with PTSD. We explicitly investigated *stable networks*, which could be summarily dismissed as negative findings since the contributing regions have minimal between-group differences. However, negative findings do not necessarily indicate that group differences in SC are absent. Negative findings may indicate insufficient statistical power. The sample size of the present study provides sufficient power to detect extremely small effect sizes, which we may confidently interpret as negative findings that reflect networks of *stable* regions.

It is important to contrast the interpretation of CT- with SA-based networks. The relationship between CT and SA is complex involving myriad factors including brain hemisphere, brain region, age, IQ, disease, genetics, and many other factors (33, 53). The large size of the human cortex, in comparison to other animals, is driven primarily by expansion of SA, not an increased CT (54), and achieved through gyral folding. Individual differences in cortical volume are largely attributable to variability in surface area as opposed to cortical thickness (55). While CT and SA are highly heritable ($r_g = 0.81$ and 0.89 , respectively), the genetic correlation between CT and SA is exceedingly low ($r_g = 0.08$). The influence of environment on CT and SA is also relatively low, accounting for 20% of their variance (56). Findings from structural MRI of 51,665 genotyped individuals show that common genetic variants explain greater phenotypic variance in SA (8 to 31%) than in CT (1 to 13%). Strikingly, 175 unique genetic loci were associated with SA, but only 10 unique loci were associated with CT (57). Understanding the functional roles of

these genetic loci will contribute to interpretation of CT-based and SA-based structural connectivity, which will help us to understand the genetic contribution of remodeling of cortical topography in PTSD. Perhaps identifying common genetic variants that explain CT- and SA-based structural connectivity between regions and within networks will provide insights into the genetic architecture of the structural connectome (10).

Patients with depression alone showed higher mean SC in both CT-based and SA-based atrophic networks, and lower mean SC in SA-based hypertrophic networks, as compared to healthy controls. These results suggest that depression is associated with more coordinated propagation of CT and SA reductions, and less coordinated SA increases. Our result is consistent with previous reports that depression is associated with widely distributed CT reductions (58). Patients with PTSD alone showed lower mean SC in CT-based atrophic networks than patients with depression alone, suggesting that PTSD is associated with more coordinated decline throughout CT-based networks than depression. We also found that PTSD with comorbid depression was associated with lower mean SC in CT-based atrophic networks than depression alone, lower mean SC in SA-based atrophic networks compared to PTSD alone and depression alone, and higher mean SC in SA-based hypertrophic networks relative to depression alone. Previous studies have documented greater volume reductions in cortical structures including anterior/middle cingulate cortex, orbitofrontal cortex, and dorsolateral prefrontal cortex in PTSD with comorbid depression that are absent in either disorder alone (59). Behaviorally, higher levels of distress (60), impaired neurocognitive function (61), and greater risk for suicide (62) are present in comorbid PTSD and depression compared to PTSD alone. PTSD with comorbid depression, relative to either disorder alone, may be associated with larger disruptions of individual cortical regions and their network SC, which may explain greater symptom severity.

We explored the modulation of PTSD-related differences in SCN by sex and age, and modulation of SCNs by PTSD symptom severity. We found that (**Supplementary Fig. S1**), females with PTSD and males without PTSD had greater SC in CT-based atrophic networks than females without PTSD. Males without PTSD had greater mean SC in CT-based stable networks than males with PTSD and females without PTSD. Diffusion-based structural connectome studies in youth show that males have stronger connections between regions for perception and coordinated action, whereas females have stronger connections between analytical and intuitive processing modes (63), demonstrating the sex-related differences in brain connections. We also found (**Supplementary Fig. S2**) an inverted U-shaped relationship between age and SC in CT-based atrophic networks that peaked at 20-30 years in non-PTSD and 15-20 years in PTSD, whereas SA-based hypertrophic networks peaked at 10-15 years in both groups. We found significant PTSD-related SC differences in some age groups, particularly < 10 years, as demonstrated by higher SC in CT-based hypertrophic and stable networks, lower SC in CT-based atrophic networks, and lower SC in SA-based hypertrophic networks. Our results suggest that multiple networks undergo transformation in a coordinated fashion to support the development of the brain as well as PTSD symptoms, particularly during early childhood. A previous longitudinal study in healthy young people (9) showed that similar global and nodal topological properties as well as mesoscopic features are shared by SC networks and maturation networks, which are based on each region's slope of maturation with age and pairwise correlations in the rate of maturation across subjects.

Strengths and Limitations

A major strength of our study is a large cohort of over 3,400 participants that represent diverse geography, demography (sex, age, race), trauma type (military, sexual violence, natural disasters) and clinical comorbidity. This sample heterogeneity enhances the generalizability and

reproducibility of our findings. Harmonization of CT and SA measures sourced from 29 international sites with different MRI scanners was addressed with *ComBat* (35). A major strength our methodology is empirical confirmation that the most atrophic regions, or most hypertrophic regions, constitute the networks with the greatest change in SC. The possibility that SC might be most affected by PTSD in networks formed of random regions, i.e. where PTSD associated changes of individual regions are completely unremarkable, has been robustly addressed.

The following limitations warrant consideration when interpreting the present results. Firstly, our study is based on cross-sectional data which lacks longitudinal information to inform neurodevelopmental processes. Combining neuroimaging data from multiple longitudinal scans on each subject over several years of follow-up, preferably with pre-trauma and post-trauma observations, may help us to better understand the developmental changes in SC networks among trauma-exposed and PTSD subjects. Secondly, image quality reflected by the Euler number was not significantly different between PTSD and non-PTSD groups in most sites except for Duke University (DeBellis) and INTRUST. Higher image quality is associated with greater CT in dorsolateral prefrontal cortex, superior parietal cortex, and lateral temporal cortex, as well as smaller CT in occipital and posterior cingulate cortex (64). Cortical morphometry and therefore SC may be biased by the PTSD-related differences in image quality at two sites. However, our leave-three-sites-out analyses indicated that our results are reliable. Future studies on cortical morphometry and cortico-cortical SCNs should consider including the image quality as a covariate in statistical models. Finally, information on illness chronicity, developmental timing of trauma, childhood maltreatment, and other comorbidities such as anxiety, were unavailable in the datasets shared with us by our Consortium partners. Future research comparing trauma-exposed individuals without PTSD to trauma-unexposed individuals could offer evidence supporting a hypothetical *resilience network*. Similarly, differences in

patients with remitted PTSD compared to chronic PTSD could support the existence of a hypothetical *recovery network*. Future research could also compare patient groups exhibiting specific symptom clusters of PTSD.

Conclusions

Cortico-cortical connections shape the topography of PTSD-related differences in cortical morphometry. Thus, regional cortical morphometry associated with PTSD, does not occur in isolated brain regions and independent of differences seen in other cortical regions. Rather, the regions whose morphometry are most affected by PTSD, albeit not significantly, form networks whose covariance structure is significantly affected by PTSD diagnosis and symptom severity. This finding fundamentally and significantly extends our understanding about the effects of PTSD on brain structure. Namely, cortical regions must be viewed from a wholistic standpoint as acting within the context of networks that are affected in coordinated manner by PTSD and further modulated by comorbid depression, sex, and age. The structural covariance networks that are perturbed in PTSD comport with converging evidence from resting state functional connectivity networks and networks impacted by stress hormones, inflammation, and epigenetics.

Acknowledgments

DoD W81XWH-10-1-0925; Center for Brain and Behavior Research Pilot Grant; South Dakota Governor's Research Center Grant; CX001600 VA CDA; NHMRC Program Grant #1073041; R01 MH111671; VISN6 MIRECC; German Research Foundation grant to J. K. Daniels (DA 1222/4-1 and WA 1539/8-2); VA RR&D 1K2RX000709; NIMH R01-MH043454; NIMH T32-MH018931; 5U01AA021681-08; K24MH71434; K24 DA028773; R01 MH63407; R01 AA12479; R01 MH61744; K99NS096116; VA RR&D 1K1RX002325; VA RR&D 1K2RX002922; MH101380; ZonMw, the Netherlands organization for Health Research and Development grant to Miranda Olf (40-00812-98-10041); Academic Medical Center Research Council grant to Miranda Olf (110614); VA CSR&D 1K2CX001680; VISN17 Center of Excellence pilot funding; NIMH R01MH105535; NIMH 1R21MH102634; German Federal Ministry of Education and Research (BMBF RELEASE 01KR1303A); German Research Society (Deutsche Forschungsgemeinschaft, DFG; SFB/TRR 58: C06, C07); R01MH111671; R01MH117601; R01AG059874; MJFF 14848; MH098212; MH071537; M01RR00039; UL1TR000454; HD071982; HD085850; R21MH112956; Anonymous Women's Health Fund; Kasparian Fund; Trauma Scholars Fund; Barlow Family Fund; W81XWH-08-2-0159; Department of Veterans Affairs via support for the National Center for PTSD; NIAAA via its support for (P50) Center for the Translational Neuroscience of Alcohol; NCATS via its support of (CTSA) Yale Center for Clinical Investigation; NIH R01 MH106574; F32MH109274; NIMH 1R21MH102634; R01MH113574; R01-MH103291; BOF 2-4 year project to Sven C. Mueller (01J05415); R01MH105355; Dana Foundation (to Dr. Nitschke); the University of Wisconsin Institute for Clinical and Translational Research; a National Science Foundation Graduate Research Fellowship (to Dr. Grupe); the National Institute of Mental Health (NIMH) R01 MH63407 (to De Bellis), R01 AA12479 (to De Bellis), and R01 MH61744 (to De Bellis); R01-MH043454 and T32-MH018931 (to Dr. Davidson); core grant to the Waisman Center from the National Institute of Child Health and Human Development (P30-HD003352); NIMH K23MH112873; Veterans Affairs Merit Review Program (10/01/08 – 09/30/13); L30 MH114379; German Federal Ministry of Education and Research (BMBF RELEASE 01KR1303A); South African Medical Research Council "SHARED ROOTS" Flagship Project; Grant MRC-RFA-FSP-01-2013/SHARED ROOTS; South African Research Chair in PTSD from the Department of Science and Technology and the National Research Foundation; US Department of Defence Grant W81XWH08-2-0159 (PI: Stein, Murray B); VA RR&D I01RX000622; CDMRP W81XWH-08-2-0038; South African Medical Research Council; NARSAD Young Investigator; K01 MH118428; Department of Defense award number W81XWH-12-2-0012; ENIGMA was also supported in part by NIH U54 EB020403 from the Big Data to Knowledge (BD2K) program; R56AG058854; R01MH116147; R01MH111671; P41 EB015922; 1R01MH110483; 1R21 MH098198; R01MH105355-01A. The views expressed in this article are those of the authors and do not necessarily reflect the position or policy of the Department of Veterans Affairs, the United States Government, or any other funding sources listed here.

Conflicts of Interest

Dr. Abdallah has served as a consultant, speaker and/or on advisory boards for FSV7, Lundbeck, Psilocybin Labs, Genentech and Janssen, and editor of Chronic Stress for Sage Publications, Inc.; he has filed a patent for using mTOR inhibitors to augment the effects of antidepressants (filed on August 20, 2018). Dr. Davidson is the founder and president of, and serves on the board of directors for, the non-profit organization Healthy Minds Innovations, Inc. Dr. Jahanshad received partial research support from Biogen, Inc. (Boston, USA) for research unrelated to the content of this manuscript. Dr. Krystal is a consultant for AbbVie, Inc., Amgen, Astellas Pharma Global Development, Inc., AstraZeneca Pharmaceuticals, Biomedisyn Corporation, Bristol-Myers Squibb, Eli Lilly and Company, Euthymics Bioscience, Inc., Neurovance, Inc., FORUM Pharmaceuticals, Janssen Research & Development, Lundbeck Research USA, Novartis Pharma AG, Otsuka America Pharmaceutical, Inc., Sage Therapeutics, Inc., Sunovion Pharmaceuticals, Inc., and Takeda Industries; is on the Scientific Advisory Board for Lohocla Research Corporation, Mnemosyne Pharmaceuticals, Inc., Naurex, Inc., and Pfizer; is a stockholder in Biohaven Pharmaceuticals; holds stock options in Mnemosyne Pharmaceuticals, Inc.; holds patents for Dopamine and Noradrenergic Reuptake Inhibitors in Treatment of Schizophrenia, US Patent No. 5,447,948 (issued September 5, 1995), and Glutamate Modulating Agents in the Treatment of Mental Disorders, U.S. Patent No. 8,778,979 (issued July 15, 2014); and filed a patent for Intranasal Administration of Ketamine to Treat Depression. U.S. Application No. 14/197,767 (filed on March 5, 2014); US application or Patent Cooperation Treaty international application No. 14/306,382 (filed on June 17, 2014). Filed a patent for using mTOR inhibitors to augment the effects of antidepressants (filed on August 20, 2018). Dr. Schmahl is consultant for Boehringer Ingelheim International GmbH. Dr. Stein has received research grants and/or consultancy honoraria from Lundbeck and Sun. Dr. Thompson received partial research support from Biogen, Inc. (Boston, USA) for research unrelated to the topic of this manuscript. All other authors report no biomedical financial interests or potential conflicts of interest.

References

1. Shalev A, Liberzon I, Marmar C (2017): Post-Traumatic Stress Disorder. *New Engl J Med.* 376:2459-2469.
2. Lindemer ER, Salat DH, Leritz EC, McGlinchey RE, Milberg WP (2013): Reduced cortical thickness with increased lifetime burden of PTSD in OEF/OIF Veterans and the impact of comorbid TBI. *Neuroimage-Clin.* 2:601-611.
3. Mueller SG, Ng P, Neylan T, Mackin S, Wolkowitz O, Mellon S, et al. (2015): Evidence for disrupted gray matter structural connectivity in posttraumatic stress disorder. *Psychiat Res-Neuroim.* 234:194-201.
4. Wrocklage KM, Averill LA, Cobb Scott J, Averill CL, Schweinsburg B, Trejo M, et al. (2017): Cortical thickness reduction in combat exposed U.S. veterans with and without PTSD. *Eur Neuropsychopharmacol.* 27:515-525.
5. Li SG, Huang XQ, Li LJ, Du F, Li J, Bi F, et al. (2016): Posttraumatic Stress Disorder: Structural Characterization with 3-T MR Imaging. *Radiology.* 280:537-544.
6. Hu H, Sun YW, Su SS, Wang Y, Qiu YM, Yang X, et al. (2018): Cortical surface area reduction in identification of subjects at high risk for post-traumatic stress disorder: A pilot study. *Aust Nz J Psychiat.* 52:1084-1091.
7. Klabunde M, Weems CF, Raman M, Carrion VG (2017): The moderating effects of sex on insula subdivision structure in youth with posttraumatic stress symptoms. *Depress Anxiety.* 34:51-58.
8. Gong GL, He Y, Chen ZJ, Evans AC (2012): Convergence and divergence of thickness correlations with diffusion connections across the human cerebral cortex. *Neuroimage.* 59:1239-1248.
9. Alexander-Bloch A, Giedd JN, Bullmore ET (2013): Imaging structural co-variance between human brain regions. *Nat Rev Neurosci.* 14:322-336.

10. Romero-Garcia R, Whitaker KJ, Vasa F, Seidlitz J, Shinn M, Fonagy P, et al. (2018): Structural covariance networks are coupled to expression of genes enriched in supragranular layers of the human cortex. *Neuroimage*. 171:256-267.
11. Sun D, Davis SL, Haswell CC, Swanson CA, LaBar KS, Fairbank JA, et al. (2018): Brain Structural Covariance Network Topology in Remitted Posttraumatic Stress Disorder. *Frontiers in Psychiatry*. 9.
12. Sun DL, Haswell CC, Morey RA, de Bellis MD (2019): Brain structural covariance network centrality in maltreated youth with PTSD and in maltreated youth resilient to PTSD. *Dev Psychopathol*. 31:557-571.
13. Sun DL, Peveril MR, Swanson CS, McLaughlin KA, Morey RA (2018): Structural covariance network centrality in maltreated youth with posttraumatic stress disorder. *J Psychiatr Res*. 98:70-77.
14. Yun JY, Boedhoe PSW, Vriend C, Jahanshad N, Abe Y, Ameis SH, et al. (2020): Brain structural covariance networks in obsessive-compulsive disorder: a graph analysis from the ENIGMA Consortium. *Brain*. 143:684-700.
15. Cauda F, Nani A, Manuello J, Premi E, Palermo S, Tatu K, et al. (2018): Brain structural alterations are distributed following functional, anatomic and genetic connectivity. *Brain*. 141:3211-3232.
16. Roos A, Fouche JP, Stein DJ (2017): Brain network connectivity in women exposed to intimate partner violence: a graph theory analysis study. *Brain Imaging Behav*. 11:1629-1639.
17. Greenberg MS, Tanev K, Marin MF, Pitman RK (2014): Stress, PTSD, and dementia. *Alzheimers Dement*. 10:S155-S165.
18. Sherin JE, Nemeroff CB (2011): Post-traumatic stress disorder: the neurobiological impact of psychological trauma. *Dialogues Clin Neurosci*. 13:263-278.

19. Katrinli S, Stevens J, Wani AH, Lori A, Kilaru V, van Rooij SJH, et al. (2020): Evaluating the impact of trauma and PTSD on epigenetic prediction of lifespan and neural integrity. *Neuropsychopharmacol.* 45:1609-1616.
20. Wolf EJ, Chen CD, Zhao X, Zhou ZW, Morrison FG, Daskalakis NP, et al. (2021): Klotho, PTSD, and advanced epigenetic age in cortical tissue. *Neuropsychopharmacol.* 46:721-730.
21. Miller MW, Lin AP, Wolf EJ, Miller DR (2018): Oxidative Stress, Inflammation, and Neuroprogression in Chronic PTSD. *Harvard Rev Psychiat.* 26:57-69.
22. Mehta ND, Stevens JS, Li ZH, Gillespie CF, Fani N, Michopoulos V, et al. (2020): Inflammation, reward circuitry and symptoms of anhedonia and PTSD in trauma-exposed women. *Soc Cogn Affect Neur.* 15:1046-1055.
23. Clausen AN, Fercho KA, Monsour M, Disner S, Salminen L, Haswell CC, et al. (2021): Assessment of brain age in posttraumatic stress disorder: Findings from the ENIGMA PTSD and brain age working groups. *Brain and Behavior.*
24. Liao W, Zhang Z, Mantini D, Xu Q, Wang Z, Chen G, et al. (2013): Relationship between large-scale functional and structural covariance networks in idiopathic generalized epilepsy. *Brain Connect.* 3:240-254.
25. Zielinski BA, Gennatas ED, Zhou JA, Seeley WW (2010): Network-level structural covariance in the developing brain. *P Natl Acad Sci USA.* 107:18191-18196.
26. Wannan CMJ, Cropley VL, Chakravarty MM, Bousman C, Ganella EP, Bruggemann JM, et al. (2019): Evidence for Network-Based Cortical Thickness Reductions in Schizophrenia. *Am J Psychiat.* 176:552-563.
27. Rakic P (1988): Specification of Cerebral Cortical Areas. *Science.* 241:170-176.
28. Rakic P (2009): Evolution of the neocortex: a perspective from developmental biology. *Nat Rev Neurosci.* 10:724-735.
29. Horton JC, Adams DL (2005): The cortical column: a structure without a function. *Philos Trans R Soc Lond B Biol Sci.* 360:837-862.

30. Sanabria-Diaz G, Melie-Garcia L, Iturria-Medina Y, Aleman-Gomez Y, Hernandez-Gonzalez G, Valdes-Urrutia L, et al. (2010): Surface area and cortical thickness descriptors reveal different attributes of the structural human brain networks. *Neuroimage*. 50:1497-1510.
31. Yang JJ, Kwon H, Lee JM (2016): Complementary Characteristics of Correlation Patterns in Morphometric Correlation Networks of Cortical Thickness, Surface Area, and Gray Matter Volume. *Sci Rep-Uk*. 6.
32. Flory JD, Yehuda R (2015): Comorbidity between post-traumatic stress disorder and major depressive disorder: alternative explanations and treatment considerations. *Dialogues Clin Neuro*. 17:141-150.
33. Schnack HG, van Haren NEM, Brouwer RM, Evans A, Durston S, Boomsma DI, et al. (2015): Changes in Thickness and Surface Area of the Human Cortex and Their Relationship with Intelligence. *Cereb Cortex*. 25:1608-1617.
34. Wierenga LM, Langen M, Oranje B, Durston S (2014): Unique developmental trajectories of cortical thickness and surface area. *Neuroimage*. 87:120-126.
35. Fortin JP, Cullen N, Sheline YI, Taylor WD, Aselcioglu I, Cook PA, et al. (2018): Harmonization of cortical thickness measurements across scanners and sites. *Neuroimage*. 167:104-120.
36. He Y, Chen ZJ, Evans AC (2007): Small-world anatomical networks in the human brain revealed by cortical thickness from MRI. *Cereb Cortex*. 17:2407-2419.
37. Benjamini Y, Hochberg Y (1995): Controlling the False Discovery Rate - a Practical and Powerful Approach to Multiple Testing. *J R Stat Soc B*. 57:289-300.
38. Crossley NA, Mechelli A, Scott J, Carletti F, Fox PT, McGuire P, et al. (2014): The hubs of the human connectome are generally implicated in the anatomy of brain disorders. *Brain*. 137:2382-2395.

39. Shang J, Lui S, Meng Y, Zhu H, Qiu C, Gong Q, et al. (2014): Alterations in low-level perceptual networks related to clinical severity in PTSD after an earthquake: a resting-state fMRI study. *PLoS One*. 9:e96834.
40. Koch SB, van Zuiden M, Nawijn L, Frijling JL, Veltman DJ, Olff M (2016): Aberrant Resting-State Brain Activity in Posttraumatic Stress Disorder: A Meta-Analysis and Systematic Review. *Depress Anxiety*. 33:592-605.
41. Ke J, Zhang L, Qi R, Xu Q, Zhong Y, Liu T, et al. (2018): Typhoon-Related Post-Traumatic Stress Disorder and Trauma Might Lead to Functional Integration Abnormalities in Intra- and Inter-Resting State Networks: a Resting-State Fmri Independent Component Analysis. *Cell Physiol Biochem*. 48:99-110.
42. Suo X, Lei D, Li K, Chen F, Li F, Li L, et al. (2015): Disrupted brain network topology in pediatric posttraumatic stress disorder: A resting-state fMRI study. *Hum Brain Mapp*. 36:3677-3686.
43. Ross MC, Cisler JM (2020): Altered large-scale functional brain organization in posttraumatic stress disorder: A comprehensive review of univariate and network-level neurocircuitry models of PTSD. *Neuroimage-Clin*. 27.
44. Michopoulos V, Powers A, Gillespie CF, Ressler KJ, Jovanovic T (2017): Inflammation in Fear- and Anxiety-Based Disorders: PTSD, GAD, and Beyond. *Neuropsychopharmacol*. 42:254-270.
45. Wellman CL, Bollinger JL, Moench KM (2020): Effects of stress on the structure and function of the medial prefrontal cortex: Insights from animal models. *Int Rev Neurobiol*. 150:129-153.
46. Godar SC, Bortolato M, Richards SE, Li FG, Chen K, Wellman CL, et al. (2015): Monoamine Oxidase A is Required for Rapid Dendritic Remodeling in Response to Stress. *Int J Neuropsychopharmacol*. 18.

47. Liston C, McEwen BS, Casey BJ (2009): Psychosocial stress reversibly disrupts prefrontal processing and attentional control. *Proc Natl Acad Sci U S A*. 106:912-917.
48. Yeshurun S, Hannan AJ (2019): Transgenerational epigenetic influences of paternal environmental exposures on brain function and predisposition to psychiatric disorders. *Mol Psychiatry*. 24:536-548.
49. Vukojevic V, Kolassa IT, Fastenrath M, Gschwind L, Spalek K, Milnik A, et al. (2014): Epigenetic modification of the glucocorticoid receptor gene is linked to traumatic memory and post-traumatic stress disorder risk in genocide survivors. *J Neurosci*. 34:10274-10284.
50. Zannas AS, Provencal N, Binder EB (2015): Epigenetics of Posttraumatic Stress Disorder: Current Evidence, Challenges, and Future Directions. *Biol Psychiatry*. 78:327-335.
51. Tozzi L, Farrell C, Booij L, Doolin K, Nemoda Z, Szyf M, et al. (2018): Epigenetic Changes of FKBP5 as a Link Connecting Genetic and Environmental Risk Factors with Structural and Functional Brain Changes in Major Depression. *Neuropsychopharmacol*. 43:1138-1145.
52. Sun D, Peverill MR, Swanson CS, McLaughlin KA, Morey RA (2018): Structural covariance network centrality in maltreated youth with posttraumatic stress disorder. *J Psychiatr Res*. 98:70-77.
53. Lyall AE, Shi F, Geng X, Woolson S, Li G, Wang L, et al. (2015): Dynamic Development of Regional Cortical Thickness and Surface Area in Early Childhood. *Cereb Cortex*. 25:2204-2212.
54. Geschwind DH, Rakic P (2013): Cortical evolution: judge the brain by its cover. *Neuron*. 80:633-647.
55. Im K, Lee JM, Lyttelton O, Kim SH, Evans AC, Kim SI (2008): Brain size and cortical structure in the adult human brain. *Cereb Cortex*. 18:2181-2191.

56. Panizzon MS, Fennema-Notestine C, Eyer LT, Jernigan TL, Prom-Wormley E, Neale M, et al. (2009): Distinct Genetic Influences on Cortical Surface Area and Cortical Thickness. *Cereb Cortex*. 19:2728-2735.
57. Grasby KL, Jahanshad N, Painter JN, Colodro-Conde L, Bralten J, Hibar DP, et al. (2020): The genetic architecture of the human cerebral cortex. *Science*. 367:1340-+.
58. Suh JS, Schneider MA, Minuzzi L, MacQueen GM, Strother SC, Kennedy SH, et al. (2019): Cortical thickness in major depressive disorder: A systematic review and meta-analysis. *Prog Neuro-Psychoph*. 88:287-302.
59. Kroes MCW, Rugg MD, Whalley MG, Brewin CR (2011): Structural brain abnormalities common to posttraumatic stress disorder and depression. *J Psychiatr Neurosci*. 36:256-265.
60. Campbell DG, Felker BL, Liu CF, Yano EM, Kirchner JE, Chan D, et al. (2007): Prevalence of depression-PTSD comorbidity: Implications for clinical practice guidelines and primary care-based interventions. *J Gen Intern Med*. 22:711-718.
61. Nijdam MJ, Gersons BPR, Olf M (2013): The role of major depression in neurocognitive functioning in patients with posttraumatic stress disorder. *Eur J Psychotraumatol*. 4.
62. Ramsawh HJ, Fullerton CS, Mash HBH, Ng THH, Kessler RC, Stein MB, et al. (2014): Risk for suicidal behaviors associated with PTSD, depression, and their comorbidity in the US Army. *J Affect Disorders*. 161:116-122.
63. Ingalhalikar M, Smith A, Parker D, Satterthwaite TD, Elliott MA, Ruparel K, et al. (2014): Sex differences in the structural connectome of the human brain. *P Natl Acad Sci USA*. 111:823-828.
64. Rosen AFG, Roalf DR, Ruparel K, Blake J, Seelaus K, Villa LP, et al. (2018): Quantitative assessment of structural image quality. *Neuroimage*. 169:407-418.

Figure Legends

Figure 1. Analyses pipelines. (A) Anatomical neuroimaging data from 29 research sites was aggregated by the ENIGMA PGC PTSD working group. Regional estimates of cortical thickness (CT) and surface area (SA) extracted from 148 cortical regions based on the Destrieux atlas (Destrieux, Fischl, Dale, & Halgren, 2010) were harmonized to remove site effects with *ComBat* approach and entered into a linear model to adjust for effects of age, age², sex, and whole-brain mean CT (or SA). The residuals were used to compute Pearson correlation coefficients for each pair of cortical regions across subjects within groups. The correlation coefficients were *r*-to-*z* transformed to improve normality and yielded a structural covariance (SC) matrix for each participant group. The cortical regions were rank ordered according to the magnitude of effect size when contrasting CT (or SA) between PTSD and non-PTSD groups. The top-*n* (*n* = 2 to 148) regions with the largest effect size of differences for PTSD > non-PTSD constituted atrophic networks, PTSD < non-PTSD constituted hypertrophic networks, while the smallest effect size stable networks. The mean SC of a given *n*-region network measured by the mean of positive correlations between all possible pairs of regions were compared to 5,000 randomly generated *n*-region networks matched for hemisphere and distance. Both global and individual tests were employed to compute statistical significance based on the proportion of mean SC values from randomly chosen sets of *n* regions that exceeded or equaled the mean SC of the actual top-*n* network. As illustrated in (B), the top-*n* (*n* = 5, 10, and 20) regions showed (i) the largest effect size in CT (or SA) for PTSD < non-PTSD (atrophic networks); (ii) the largest effect size of PTSD > non-PTSD (hypertrophic networks); or (iii) the smallest effect size of PTSD vs. non-PTSD (stable networks). (C) CT-based hypertrophic networks for top-3, top-10 and top-50 regions.

Figure 2. The top-20 regions showing PTSD-related differences. The top-20 regions that (A) PTSD < non-PTSD and (B) PTSD > non-PTSD in cortical thickness. The top-20 regions that (C)

PTSD < non-PTSD and (D) PTSD > non-PTSD in surface area. Node size represents the magnitude of effect size for between-group differences per region. Warm color denotes PTSD > non-PTSD, and cool color denotes PTSD < non-PTSD. Regions names are listed in **Supplementary Table S4**. Two examples are shown on the right to denote the node size and the corresponding effect size (Cohen's d). The directions of the brain maps (axial view) are also shown.

Figure 3. Mean SC of patients with PTSD. Global tests showed that PTSD patients have higher mean SC in both CT- ($p < 0.001$) and SA-based ($p = 0.017$) atrophic networks, both CT- ($p = 0.029$) and SA-based ($p = 0.017$) hypertrophic networks, and CT-based ($p < 0.001$) but not SA-based ($p > 0.5$) stable networks than the corresponding random networks. The curves of networks with up to 50 nodes are shown for illustrative purposes, given that the mean SC of actual networks and the mean SC of the average of random networks were very similar for large network sizes. Red curve, mean SC of the actual networks; Blue curve, mean SC of the average of 5,000 random networks; light blue ribbon, 95% confidence interval (CI) of the 5,000 random networks.

Figure 4. Mean SC of trauma-exposed participants without PTSD. Global tests showed that participants without PTSD had higher mean SC in both CT- ($p < 0.001$) and SA-based ($p < 0.001$) atrophic networks, SA-based ($p = 0.014$) but not CT-based ($p = 0.139$) hypertrophic networks, and neither CT- ($p = 0.264$) nor SA-based ($p = 0.732$) stable networks than in corresponding random networks. The curves for networks with up to 50 nodes are shown for illustrative purpose, given that the mean SC of actual networks and the mean SC of the average of random networks were very similar for large network sizes. Red curve, mean SC of the actual networks; Blue curve, mean SC of the average of 5,000 random networks; light blue ribbon, 95% confidence interval (CI) of the 5,000 random networks.

Figure 5. Mean SC of PTSD vs. non-PTSD. Global tests showed that patients with PTSD versus non-PTSD participants had lower mean SC in both CT- ($p = 0.014$) and SA-based ($p = 0.024$) atrophic networks, but no significant difference in CT- ($p = 0.098$) and SA-based ($p > 0.5$) hypertrophic networks as well as CT- ($p > 0.5$) and SA-based ($p > 0.5$) stable networks. The curves of networks with up to 50 nodes are shown for illustrative purpose, given that the mean SC of actual networks and the mean SC of the average of random networks were very similar for large network sizes. Red curve, mean SC of the actual networks; Blue curve, mean SC of the average of 5,000 random networks; light blue ribbon, 95% confidence interval (CI) of the 5,000 random networks.

Figure 6. Replication analyses results. The global-tests results shown in **Figures 3, 4, and 5** are reliable as underscored by the area under curve (AUC) of mean SC for the results based on all 29 sites (represented by the red vertical line) was always located within the 95% confidence interval (represented by two blue vertical dashed lines) of the AUC of mean SC from 5,000 iterations leaving out 3 sites at each iteration across all types of networks.

Figure 7. Interaction effects of PTSD and depression. Global tests showed that patients with depression alone had higher mean SC in **(A)** CT-based ($p < 0.001$) and **(B)** SA-based ($p < 0.001$) atrophic networks, and lower mean SC in **(D)** SA-based hypertrophic networks ($p = 0.029$), than patients with both PTSD and depression. Patients with depression alone also showed higher mean SC in both **(A)** CT-based ($p < 0.001$) and **(B)** SA-based ($p < 0.001$) atrophic networks, and lower mean SC in **(D)** SA-based hypertrophic networks ($p < 0.001$), than patients with neither PTSD nor depression. Patients with PTSD alone showed lower mean SC in **(A)** CT-based atrophic networks than patients with depression alone ($p < 0.001$), and higher mean SC in **(B)** SA-based atrophic networks than patients with both PTSD and depression ($p < 0.001$) as well as participants with neither PTSD nor depression ($p = 0.014$). No significant PTSD x depression interaction effect (global p -values > 0.2) was found in the other types of

networks shown in (C), (E) and (F). The curves of networks with up to 30 nodes were shown for illustrative purposes. Error bar denotes 95% confidence interval of 5,000 random networks. * represents $p < 0.05$; *** represents $p < 0.001$.

Table 1. Demographic and clinical information per site.

Site	Number of Participants				Age (years)	Trauma	MDD (%)	Type
	CT	SA	Male/ Female	PTSD/ non-PTSD				
ADNIDOD	194	194	193/1	80/106	69.0±5	Y	2.5	Military
Booster (AMC)	75	75	40/35	38/37	40.0±10.0	Y	31	Police
Columbia	88	88	31/57	53/35	36.0±9.8	Y	24	Civilian
Duke University (DeBellis)	115	117	53/62	29/86	10.0±2.6	Y/N	N/A	Civilian
Minneapolis VAMC	169	171	161/8	74/95	33.0±7.9	Y	28.4	Military
Duke University /Durham VA	385	385	310/75	114/270	40.0±10.0	Y	40.3	Both
Ghent	67	67	0/67	8/59	37.0±12.0	N	46.3	Civilian
Groningen (Charité Berlin)	40	40	0/40	40/0	38.0±10.0	Y	67.5	Civilian
University of Wisconsin (Grupe)	57	58	53/4	19/38	31.0±6.4.0	Y	100	Military
Emory GTP	174	174	5/169	66/108	38.0±13.0	Y	51.7	Civilian
INTRUST	373	373	220/145	109/262	35.0±14.0	Y	21.7	Both
University of Wisconsin (Larson)	67	67	33/34	20/47	33.0±11.0	Y	0	Civilian
Leiden	52	52	7/45	22/30	15.2±2.0	N	19.2	Civilian
Mannheim	48	48	0/48	48/0	36.0±12.0	Y	97.9	Civilian
McLean	52	52	0/52	39/13	38.0±12.0	Y	75	Civilian
Muenster	47	47	5/42	21/26	27.0±7.0	Y	34	Civilian
Phan	43	43	43/0	23/20	32.0±8.0	Y	53.5	Military
McLean (Rosso)	106	97	49/57	21/85	34.0±9.0	Y	23	Civilian
University of Toledo	76	76	42/34	15/61	35.0±11.3	Y	41	Both
UCAS	70	70	32/38	34/36	50.0±7.0	Y	64.3	Civilian
Cape Town	62	63	0/62	7/55	29.0±8.0	Y	50	Civilian
University of Washington	255	255	125/130	53/202	14.0±3.1	Y	15.3	Civilian
WACO VA	66	66	56/10	41/25	41.0±11.1	N	67	Military
WestHaven VA	72	71	63/8	34/40	35.0±10.0	Y	75	Military
Yale	70	70	59/11	22/48	29.2±9.2	Y	0	Civilian




UNSW	162	163	63/99	49/113	40.4±8.0	Y	28.4	Civilian
South Dakota	123	123	99/24	78/45	29.0±7.0	Y	35	Both
Stellenbosch	260	260	72/188	121/139	41.0±13.0	Y	0	Civilian
Stanford	71	71	29/41	70/1	37.0±11.3	Y	0	Civilian
Total	3438	3436	1843/ 1586	1350/ 2076	-	-	29.9	-



Note: CT = cortical thickness; SA = surface area; Trauma = whether the non-PTSD participants are trauma-exposed; MDD (%) = percentage of participants who have major depressive disorder; Type = participants are from military/police, civilian, or both units.

Table 2. Area under curve (AUC) of mean SC for the actual network and the average of 5,000 random networks.



Network Type	CT-based networks				SA-based networks			
	Act.	Rand.	.95 CI	Global p	Act.	Rand.	.95 CI	Global p
	<u>PTSD</u>							
Atrophic	13.975	12.195	[11.918, 12.572]	<0.001***	9.480	8.725	[8.494, 9.126]	0.017*
Hypertrophic	12.846	12.104	[11.839, 12.512]	0.029*	9.356	8.692	[8.483, 9.061]	0.017*
Stable	13.211	12.193	[11.938, 12.567]	<0.001***	8.652	8.689	[8.473, 9.049]	>0.500
	<u>non-PTSD</u>							
Atrophic	14.483	12.397	[12.112, 12.785]	<0.001***	9.616	8.511	[8.286, 8.918]	<0.001***
Hypertrophic	12.832	12.317	[12.049, 12.729]	0.139	9.050	8.450	[8.242, 8.804]	0.014*
Stable	11.977	12.260	[11.983, 12.642]	0.264	8.798	8.566	[8.363, 8.890]	0.732
	<u>PTSD versus non-PTSD</u>							
Atrophic	-0.507	-0.205	[-0.382, -0.037]	0.014*	-0.136	0.211	[0.052, 0.372]	0.024*
Hypertrophic	0.015	-0.212	[-0.390, -0.037]	0.098	0.332	0.240	[0.079, 0.403]	>0.500
Stable	-0.155	-0.141	[-0.312, 0.033]	>0.500	0.172	0.215	[0.062, 0.376]	>0.500

Note: Act. = mean SC of the actual network; Rand. = average of the mean SC of 5,000 random networks; .95 CI = 95% confidence interval of the mean SC of 5,000 random networks; Global p = global p value (Bonferroni corrected) for the actual-versus-random comparison. *, $p < 0.05$; ***, $p < 0.001$.





● Cohen's $d = 0.05$
● Cohen's $d = 0.1$



Anterior
Left ← → Right
Posterior






CT-based




Atrophic

Hypertrophic

Stable




SA-based






CT-based


Atrophic






Hypertrophic



Stable




SA-based






CT-based


Atrophic






Hypertrophic



Stable



SA-based



— AUC of All-Sites Results
 - - - 95% CI of Leave-3-Sites-Out Results

CT-based

SA-based

